DATE \$1/05/96 OPI APPLN. ID 38018/95 AOJR DATE/18/07/96 PCT NUMBER PCT/DK95/00432



(51) International Patent Classification 6: C11D 1/66, 17/00

(11) International Publication Number:

WO 96/14376

A1

(43) International Publication Date:

17 May 1996 (17.05.96)

(21) International Application Number:

PCT/DK95/00432

(22) International Flling Date:

I November 1995 (01.11.95)

(30) Priority Data:

1279/94

4 November 1994 (04.11.94)

DK

(71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Alle, DK-2880 Bagsværd (DK).

(72) Inventor; and

(75) Inventor/Applicant (for US only): OLESEN, Tine [DK/DK]; Novo Nordisk a/s. Novo Allé, DK-2880 Bagsværd (DK).

(74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA. CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).

Published

With international search report.

687269

(54) Title: GLYCOLIPID/MPG GELS

(57) Abstract

This invention relates to an environmental friendly detergent composition useful in a gel for cleaning hard surfaces comprising glycolipid and polyalcohol.

1

GLYCOLIPID/MPG GELS

FIELD OF INVENTION

This invention relates to an environmental friendly detergent composition useful in a gel for cleaning hard surfaces.

BACKGROUND OF THE INVENTION

In industry hard surface cleaning has traditionally been carried out by high pressure foam cleaning. There are some severe safety and environmental problems by using high pressure foam cleaning such as high pH, aggressive cleaning agents, high water consumption, aerosol formation and mechanical noise.

To overcome these safety and environmental problems much effort has been put into developing different hard surface cleaners in the form of gels. The gel cleaners have the 15 advantages of giving no or insignificant aerosol formation, they have a low water consumption, and they are easy to apply: Prior to dilution the gel compositions are readily mobile liquids which are convenient for transport etc., but when diluted to working concentrations they start to thicken to a 20 gel-like consistency, which mean that they will remain in contact with the surface or object to be cleaned for sufficient time to enable the cleansing to occur (for reference see EP 0 314 232), so if the detergents of the gel composition are environmental friendly, the gel application method is a very safe and gentle way of cleaning hard surfaces.

The cleaning gel compositions proposed until now have contained surfactants, builders and thickeners (see Research Disclosure 34045, August 1992/619), the surfactants e.g. being benzene or naphthalene sulphate or sulphonate derivatives (see SP 0 314 232), so there is a need for "more green" detergent gel compositions.

SUMMARY OF THE INVENTION

In this invention it is surprisingly found that a gel can be made of very environmental friendly chemicals, namely glycolipid, polyalcohols and water.

Accordingly, the present invention relates to a liquid detergent composition which undergoes a viscosity increase upon aqueous dilution, comprising

- a) one or more glycolipids
- b) one or more polyalcohols
- 10 c) water.

DETAILED DISCLOSURE OF THE INVENTION

In this invention it is surprisingly found that a gel cleaning composition can be made of glycolipid(s), polyalcohol(s) and water.

According to the invention the glycolipid may be a sugar or sugar alcohol fatty acid ester (as described in JP 63-112,993) or a derivative thereof or a fatty acid monoester or a mixture of fatty acid monoesters of alkylglycoside (as described in US 5,191,071 and in US 5,200,328) or a derivative thereof. In particular the glycolipid is a fatty acid monoester of ethylglycoside. the fatty acid containing 8-22 carbon atoms, e.g. ethyl 6-0-dodecanoyl glucoside.

According to the invention any polyalcohol which has two or more hydroxyl groups may be used in making the gel. Itseful examples are 1,2-ethanediol, 1,2-propandiol, 1,3-propandiol, 1,2,3-propantriol, butanediols, butanetriols, pentanediols, and sugar alcohols. In particular 1,2-propandiol is preferred.

The gel composition may optionally also contain other 30 detergent ingredients such as solvents and sequestrants.

glucoside, 1,2-propandiol and water (in lower concentrations the g l formation decreases, for reference see Example 1).

The gel composition may also contain one or more enzymes. The enzymes are in particular proteases (for instance 5 Savinase 16.0 L, Alcalase 2.5 L, Esperase 8.0 L or Durazym 16.0 L, all available from Novo Nordisk A/S), amylases (for instance Termamyl 300 L available from Novo Nordisk A/S), lipases (for instance Lipolase 100 L available from Novo Nordisk A/S) or cellulases (for instance Celluzyme 1.0 L available from Novo Nordisk A/S). The amount of enzyme may be dosed so that the percentage of enzyme in the finished gel will be of 0.001-10%, preferably 0.01-1%, in particular about 0.01-0.5%.

Gel Making in situ

The gel is made in situ. The detergent composition comprising the glycolipid(s) and the polyalcohol(s) are diluted with water and applied to the soiled surfaces by using some Find of application system (e.g. diluted through a venturi and applied via a special lance). The detergent composition is best applied using gel generation equipment which automatically draws in the required percentage. The required percentage of a given glycolipid and a given polyalcohol may be found as described in Example 1. Application equipment is available from for instance Scanio A/S, Blytaekkervej 4-6, DK-9000 Aalborg, Benmark.

Cleaning Hard Surfaces

The gel is applied to the soiled surfaces as described above. The gel will set on the surfaces within a few seconds. The gel will remain in its form even on complicated shaped equipment, vertical surfaces and ceilings and will therefore give very long contact times. Dwell times are difficult to estimate beforehand and should be determined by trial. A dwell time of 5-30 minutes may often be required.

After a sufficient dwell time to allow for surfactant is action the gel composition is rinsed off using a washing syst m. In some cases it may be necessary to use a pressure

washing system to rinse off the gel composition, but often removal by gentle flushing with water is easy and all that is needed.

Potential Applications

The cleaning gel detergent described in this invention may be used in all kinds of industries: in dairies, in slaughterhouses, in breweries, in sea food production units etc. It may also be used in the transport sector, for instances as a cleaning agent in car washing and for general vessel wash.

10 Especially, the cleaning gel may be useful in removing oils (vegetable, animal and mineral). It may also work as a general purpose cleaner in households, e.g., for cleaning windows, bathrooms and kitchens, in particular it may be useful as an efficient oven cleaner. It may also be useful for janitorial purposes such as computer cabinet cleaning etc.

The cleaning gel of the present invention may also be very useful for personal care purposes, in skin, hair and dental applications, such as cleaning dentures.

The invention is further illustrated in the following more example which is not intended to be in any way limiting to the scope of the invention as claimed.

EXAMPLE 1

Ethyl 6-0-dodecanoyl glucoside/1.2-propandiol/water gel

Different formulations were made by mixing warm 25 (about 55°C) ethyl 6-0-dodecanoyl glucoside with 1,2-propandiol. (The temperature of the glycolipid should be chosen so it is possible to mix the glycolipid with the polyalcohol). Each formulation was then diluted with cold water in the interval of from 0 to 90% w/w, mixed on a whirler, and the rheology of each 30 gel was visually evaluated. After 5 minutes the gel was transferred to a Haake viscosimeter (Haake Rotovisco, RV 3, with the sensor system MV I, operated according to Manu-

facturer's instructions), and the visc sity was measured at several shear rates (after the bubbles had disappeared).

All measurements were made at room temperature.

The formulations tested and the achieved results are listed in 5 Table 1 (see below).

Influence of NaCl (0-8%) was also tested, by mixing with cold NaCl-solutions, instead of cold water. The results are listed in Table 2 (see below).

Results and discussion:

The formulations prepared do not exhibit newtonian behaviour, so the viscosities of different gels are only comparable when measured at the same rotation speed. The preferred ratio between ethyl 6-0-dodecanoyl glucoside and 1,2-propandiol is 80/20. Formulation no. 2 in Table 1 is clear and has an acceptable low viscosity, like a detergent, and the formulation gives good gels when diluted to a maximum of 80% water (see no. 3 to no. 7 in Table 1).

Influence of NaCl: It can be seen from Table 2 that a NaCl-concentration from about 0.2% lowers the viscosity significantly. It can also be seen that if the gels have to contain NaCl, a concentration of about 8% is needed to maintain the viscosity (compared with no added NaCl). At none of the NaCl-concentrations tested could an increase in the viscosity be measured.

						++++ = thick "ge		•																										
	Viscosity (cp) n=32 rnm	20135	C7164	2112	>4125	>4125	•	·	1134	243	>4125	1109	120	4	532	825	400	20	2112	458	87	20	404	338	462	20	998	215	28	53	80	33	128	
	Viscosity Viscosity V	5	000017	2063	>16500	10808	5198	5115	1683	363	8234	1139	132		1205	1848	528	0	2013	479	333	9 0	363	726	644	33	875	132	17	0	0	17	99	
W/W)	Visually Judgement	1777	÷ =	:	++++ +	++++	‡	++++	‡	+	*	‡	+	+	+ + +	‡	‡	+	‡	‡	+	+	‡	+	‡	+	+	+	+	+	+	+	+	
on * 1	H ₂ 0 %	6	, <u>,</u>	3 5	2	9	2	75	80	90	0	25	20	9	70	75	80	90	0	52	20	9	20	75	8	90	0	22	20	9	20	75	8	
FIONS (Based	1.2 PRO- PANDIOL %	20	<u></u>	2 5	2 6	жэ ч	٩	S	❤	2	40	30	. 50	16	12	10	&	₹	09	45	30	24	18	15	12	9	80	09	40	32	54	20	16	
"Gel"-formulations	6-0-DODECANOYL GLUCOSIDE *	80	09	(7	2 6	75	57	02	16	ထ	09	45	30	24	18	15	12	9	40	30	20	16	12	01	ထ	4	20	15	9	æ	9	ഹ	₹	
TABLE 1:	NUMBER	-	7	m	•	r u	ი (ا م	~ . (ω	σ,	2	=	12	13	7	15	16	17	18	19	20	21	22	23	24	25	56	27	88	53	30	31	

M/H)
=
*
0
(Based
MACI
2
Ī
"Gel"-formulations
"
LABLE

							10110	-) = - -		+																				
	Viscosity (cP)	3	C7164	2/4/	2970	3630		•	+ 0/51	2104		121/	1110	1205	unstable	05	S 55	unstable	813	413	722	474	474	421	122	21	12	74	33	87
2	Viscosity (cP)	10201	7707	280c	5610	6848	5115	0110	1370	4488		8777	1/99	2178	2558	132	83	1402	1351	663	1040	842	528	627	314	11	17	83	17	99
(Based on % W/W)	Visually Judgement	7777	1111	***	+ + + + + + + + + + + + + + + + + + +	++++	‡ ‡	‡	: ‡	÷ ++	-	;	‡	‡	‡	: +	+	‡	‡	+	‡	‡	‡	‡	+	+	+	+	+	+
	NaCl %	c	0 15		9.0	m	0	0.175	0.7	3.5	c	~	C. O	1	0	0.2	0.8	4	.0	4	80	0	3.75	7.5	0	0.225	4.5	0	❖	ထ
ATIONS with Maci	1.2 PRO- PANDIOL %	œ	• •	o	o (∞	9	9	9	9	12	2 2	.	12	4	♥	4	₹	80	8	80	15	15	15	2	2	2	16	9	16
"Gel"-formulations	6-0-DODECANOYL GLUCOSIDE X	32	32	3.2	, ,	35	24	24	24	24	82	<u> </u>	2 .	81	16	91	16	16	12	12	12	10	01	10	80	ထ	∞ .	₹	❖	₹
ABLE 21	NUMBER	-	7	~	· •	+	S	9	7	&	0		? :	=	12	13	14	15	16	17	18	19	20	21	22	23	24	25	5 6	27

The Claims defining the invention are as follows:

- 1. A method of making a gel in situ comprising a liquid detergent comprising incorporating into a material to be gelled
 - a) one or more glycolipids
 - b) one or more polyalcohols
 - c) water,

5

wherein the glycolipid is a sugar fatty acid ester, a derivative of a sugar fatty acid ester, a sugar alcohol fatty acid ester, a derivative of a sugar alcohol fatty acid ester, a fatty acid monoester of an alkylglycoside or a derivative of a fatty acid monoester of an alkylglycoside.

- 2. The method according to claim 1, wherein the glycolipid is a fatty acid monoester of ethylglycoside, the fatty acid containing 8-22 carbon atoms.
- 3. The method according to claim 2, wherein the glycolipid is ethyl 6-O-dodecanovl glucoside.
- The method according to any one of claims 1 to 3, wherein the polyalcohol is a diol.
 - 5. The method according to claim 4, wherein the diol is selected from the group consisting of ethanediol, propandiol, butanediol and propanediol.
 - 6. The method according to claim 5, wherein the diol is 1,2-propandiol.
- 7. A method of making a gel in situ, substantially as hereinbefore described with reference to any one of the Examples.
 - 8. A gel formed by the method of any one of claims 1 to 7.

Dated 20 May, 1997 Novo Nordisk A/S

Patent Attorneys for the Applicant/Nominated Person SPRUSON & FERGUSON



INTERNATIONAL SEARCH REPORT

International application No. PCT/DK 95/00432

A. CLAS	SIFICATION OF SUBJECT MATTER		
IPC6:	C11D 1/66, C11D 17/00 to International Patent Classification (IPC) or to both	national classification and IPC	
	DS SEARCHED		
İ	documentation searched (classification system followed	by classification symbols)	
IPC6:			in the fields manched
	ation searched other than minimum documentation to the	he exists that such documents are included	in the relative searched
	FI,NO classes as above		
Electronic (data base consulted during the international search (nam	ne of data base and, where practicable, search	in terms used)
WPI, C	Α		
C DOCI	UMENTS CONSIDERED TO BE RELEVANT	· · · · · · · · · · · · · · · · · · ·	-
Category*	Citation of document, with indication, where a	ppropriate, of the relevant passages	Relevant to claim No.
Y	EP 0499434 A1 (UNILEVER PLC ET (19.08.92), page 9, line 2 and 14, abstract	AL), 19 August 1992 - line 10, claims 1	1-5
			·
Y	WO 9213055 A1 (HENKEL KOMMANDITO AKTIEN), 6 August 1992 (06.0 line 20 - line 21; page 4, abstract	08.92), page 3,	1-5
·			
A	EP 0491532 A1 (UNILEVER PLC ET A (24.06.92)	AL), 24 June 1992	1-5
Furth	er documents are listed in the continuation of Bo	x C. X See patent family annex	.
	categories of cited documents	later document published after the upti- date and not so conflict with the appli	can on but outed to understand
to be of	particular relevance particular relevance	the practiple or theory underlying the "X" document of particular relevance: the	
"L" docume	programment but published on or after the enternational filing date and which may throw doubts on priority claim(s) or which is establish the publication date of another claims or other	considered movel or cannot be considered and the document at taken along the document at taken along the document of particular leaves and the document of particular leaves at taken along the document at taken along the docume	ared to usvolve as usvessave
special (reason (as specified) not referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance the considered to in volve an inventive fire combined with one or more other suc	p when the document is
"P" docume	or published prior to the international filing date but later than	have charged to a names shilled to \$	x ert
	nity day claimed	Date of mailing of the international	
			21.02.96
20 Febra	uary 1996	Authorized (Ticer	
	mailing address of the ISA/	Audionzai ilica	İ
	Patent Office S-102 42 STOCKHOLM	Dagmar Järvman	
	1 +46 9 666 02 96	Telephone No. +46 8 782 25 00	

INTERNATIONAL SEARCH REPORT Information in patent family members

05/02/96

International application No.
PCT/DK 95/00432

	locument arch report	Publication date	Patent family member(s)	Publication date
P-A1-	0499434	19/08/92	SE-T3- 049943 CA-A- 2060698 DE-D,T- 69203473 JP-A- 5059394 US-A- 5417879	13/08/92 11/01/96 09/03/93
D-A1-	9213055	06/08/92	DE-A- 4102502 EP-A- 0569370 JP-T- 6504553	18/11/93
-A1-	0491532	24/06/92	JP-A- 430099	23/10/92

THIS PAGE BLANK (USPTO)